

AMLODIPINE BIOTECH

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS:

S3

AMLODIPINE 5 BIOTECH tablets
AMLODIPINE 10 BIOTECH tablets
Amlodipine maleate
Sugar free

Read all of this leaflet carefully before you start taking

AMLODIPINE BIOTECH

- Keep this leaflet. You may need to read it again
- If you have further questions, please ask your doctor, pharmacist, nurse or other healthcare provider.
- AMLODIPINE BIOTECH has been prescribed for you personally and you should not share your medication with other people. It may harm them, even if their symptoms are the same as yours.

What is in this leaflet

1. What AMLODIPINE BIOTECH is and what it is used for
2. What you need to know before you take AMLODIPINE BIOTECH
3. How to take AMLODIPINE BIOTECH
4. Possible side effects
5. How to store AMLODIPINE BIOTECH
6. Contents of the pack and other information

1. What AMLODIPINE BIOTECH is and what it is used for

Amlodipine maleate, the active ingredient in AMLODIPINE 5 BIOTECH and AMLODIPINE 10 BIOTECH, is a selective calcium channel blocker that have an effect on the blood vessels.

AMLODIPINE BIOTECH is used to treat angina pectoris and mild- to moderate hypertension, alone or in combination with other antihypertensive medicines.

2. What you need to know before you take AMLODIPINE BIOTECH Do not take AMLODIPINE BIOTECH:

- if you are hypersensitive (allergic) to amlodipine maleate or any of the other ingredients of AMLODIPINE BIOTECH (listed in section 6).
- if you are hypersensitive to dihydropyridines.

Warnings and precautions

Special care should be taken with AMLODIPINE BIOTECH:

- if you are elderly and your dose needs to be increased
- if you have liver disease
- if you have heart failure
- if you had a recent heart attack

Children

Safety and efficacy have not been established.

Other medicines and AMLODIPINE BIOTECH

Always tell your healthcare provider if you are taking any other medicine. (This includes all complementary or traditional medicines.)

AMLODIPINE BIOTECH may be affected by other medicines, such as:

- ritonavir, indinavir, nelfinavir (so called protease inhibitors used to treat HIV)
- ketoconazole, itraconazole (anti-fungal medicines)
- rifampicin, erythromycin, clarithromycin (antibiotics)
- verapamil, diltiazem (heart medicines)
- hypericum perforatum (St. John's Wort)
- dantrolene (infusion for severe body temperature abnormalities)

AMLODIPINE BIOTECH may affect other medicines, such as:

- tacrolimus, sirolimus, temsirolimus, and everolimus (medicine used to alter the way your immune system works)
- ciclosporin (an immunosuppressant)
- simvastatin (a cholesterol lowering medicine)

AMLODIPINE BIOTECH with food and alcohol

Grapefruit juice and grapefruit should not be consumed by people who are taking AMLODIPINE BIOTECH. This is because grapefruit and grapefruit juice can lead to an increase in the blood levels of the active ingredient amlodipine, which can cause an unpredictable increase in the blood pressure lowering effect of amlodipine.

Pregnancy, breastfeeding and fertility

If you are pregnant or breastfeeding, think you might be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other healthcare provider for advice before taking this medicine.

Pregnancy

The safety of AMLODIPINE BIOTECH in human pregnancy has not been established.

Breastfeeding

Amlodipine has been shown to pass into breast milk in small amounts.

Driving and using machines

If you may feel nauseous, dizzy, tired or have a headache do not drive or operate machinery, especially at the beginning of taking AMLODIPINE BIOTECH.

It is not always possible to predict to what extent AMLODIPINE BIOTECH may interfere with your daily activities. You should ensure that you do not engage in driving a vehicle or use machines until you are aware of the measure to which AMLODIPINE BIOTECH affects you.

3. How to take AMLODIPINE BIOTECH

Do not share medicines prescribed for you with any other person. Always take AMLODIPINE BIOTECH exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Your doctor will tell you how long your treatment with AMLODIPINE BIOTECH will last. Do not stop treatment early. If you have the impression that the effect of AMLODIPINE BIOTECH is too strong or too weak, tell your doctor or pharmacist.

Take the tablets with a sufficient quantity of liquid (e.g., one glass of water)

The recommended dose of 5 mg AMLODIPINE BIOTECH once daily. If there is no improvement after 10 to 14 days of therapy, AMLODIPINE BIOTECH doses can be increased to 10 mg once daily.

If your take more AMLODIPINE BIOTECH than you should

Taking too many tablets may cause your blood pressure to become low or even dangerously low. You may feel dizzy, lightheaded, faint or weak. If blood pressure drop is severe enough shock can occur. Your skin could feel cool and clammy and you could lose consciousness. Excess fluid may accumulate in your lungs (pulmonary oedema) causing shortness of breath that may develop up to 24 - 48 hours after intake.

In the event of an overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison centre.

If you forget to take AMLODIPINE BIOTECH

Do not take a double dose to make up for forgotten individual doses.

4. Possible side effects

AMLODIPINE BIOTECH can have side effects.

Not all side effects reported for AMLODIPINE BIOTECH are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking AMLODIPINE BIOTECH, please consult your healthcare provider for advice.

If any of the following happens, stop taking/ using AMLODIPINE BIOTECH and tell your doctor immediately or go to the casualty department at your nearest hospital:

- swelling of the hands, feet, ankles, face, lips and mouth or throat, which may cause difficulty in swallowing or breathing,
- rash or itching,
- fainting.

These are all very serious side effects. If you have them, you may have had a serious reaction to AMLODIPINE BIOTECH. You may need urgent medical attention or hospitalisation.

Tell your doctor immediately or go to the casualty department at your nearest hospital if you notice any of the following:

- deficiency of the platelets in the blood, causing bleeding into the tissues, bruising and slow blood clotting after injury (thrombocytopenia)
 - reduction in the number of white cells in the blood (leukocytopenia)
 - excess of glucose in the bloodstream (hyperglycaemia)
 - temporary loss of consciousness caused by a fall in blood pressure
 - rapid, strong, or irregular heartbeat
 - heart attack
 - heartbeat is irregular (slow, rapid) or have an abnormal rhythm (heart's two upper chambers (the atria) beat chaotically and irregularly (atrial fibrillation))
 - difficulty in breathing
 - inflammation of the pancreas
 - inflammation of the liver
 - elevated liver enzymes and liver disease
- These are all serious side effects. You may need urgent medical attention.

Tell your doctor if you notice any of the following:

Frequent side effects:

- dizziness
- headache (especially at the beginning of the treatment)
- drowsiness
- visual disturbance and double vision
- flushing, become red and hot
- nausea
- abdominal pain
- indigestion (dyspepsia)
- diarrhoea
- constipation
- swelling of the ankles
- muscle cramps
- extreme tiredness
- physical weakness or lack of energy
- excess of watery fluid collecting in the cavities or tissues of the body (oedema)

Less frequent side effects:

- Depression
- Mood changes, including feeling of worry and nervousness
- Sleepiness
- confusion
- high levels of muscle tone or tension (hypertonia)
- loss of sensation in the body (hypoesthesia)
- pins and needles feeling (paraesthesia)
- weakness, numbness and pain, usually in your hands and feet (peripheral neuropathy)
- involuntary movement (tremor)
- disfunction in the sense of taste (dysgeusia)
- ringing or buzzing in the ears
- low blood pressure
- inflammation of the blood vessels
- coughing
- rhinitis
- vomiting
- dry mouth
- overgrowth of gum tissue around the teeth
- inflammation of the lining of the stomach
- yellowing of the skin or whites of the eyes
- complete absence of hair from areas of the body where it normally grows, baldness (alopecia)
- a rash of purple spots on the skin caused by internal bleeding from small blood vessels (purpura)
- skin discolouration
- abnormally excessive sweating that is not necessarily related to heat or exercise (hyperhidrosis)
- itching of the skin (pruritus)
- rash
- red, raised, itchy rash
- swelling around the eyes and lips (angioedema)
- skin immune reaction (erythema multiforme)
- redness and peeling of the skin (exfoliative dermatitis)
- is a rare, serious disorder of the skin and mucous membranes. It's usually a reaction to medication that starts with flu-like symptoms, followed by a painful rash that spreads and blisters. Then the top layer of affected skin dies, sheds and begins to heal after several days (Steven-Johnson syndrome).
- joint pain
- back pain
- muscle pain
- need to urinate more often than usual
- abnormalities in the process of urine voiding, including bladder control, frequency of urination, as well as the volume and composition of urine.
- wake up during the night to urinate (nocturia)
- enlargement of a man's breasts (gynaecomastia)
- inability for a man to achieve an erection or orgasm (impotence)
- pain
- chest pain
- general feeling of discomfort, illness, or unease (malaise)
- weight increase
- weight decrease

Side effects with unknown frequency:

- life-threatening skin disorder characterised by a blistering and peeling of the skin (toxic epidermal necrolysis).

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects

If you get side effects, talk to your doctor, pharmacist or nurse. You can also report side effects to SAHPRA via the **"6.04 Adverse Drug Reaction Reporting Form"**, found online under SAHPRA's publications: <https://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of AMLODIPINE BIOTECH.

5. How to store AMLODIPINE BIOTECH

Store all medicines out of reach of children.
Store at or below 25 °C.
Protect from moisture.
Return all unused medicine to your pharmacist.
Do not dispose of unused medicine in drains or sewerage systems (e.g., toilets).

6. Contents of the pack and other information

What AMLODIPINE BIOTECH contains

The active substance is amlodipine maleate equivalent to amlodipine. AMLODIPINE BIOTECH is available in two strengths, containing either 5 mg or 10 mg amlodipine.

The other ingredients are microcrystalline cellulose (Avicel PH 112), calcium hydrogen phosphate anhydrous, magnesium stearate and sodium starch glycolate (Type A) (sodium carboxymethyl starch).

What AMLODIPINE BIOTECH looks like and contents of the pack

AMLODIPINE 5 BIOTECH tablets are white to off-white, uncoated, oblong tablets.
AMLODIPINE 10 BIOTECH tablets are white to off-white, uncoated, oblong tablet, scored on one side.
AMLODIPINE BIOTECH is packed in white opaque blisters and white opaque plastic securitainers of 28, 30 and 100 tablets.
PE zip lock patient ready packs (PRPs) are available in pack sizes of 28 tablets.
All pack sizes and strengths may not necessary be marketed at one time.

Holder of Certificate of registration

Biotech Laboratories (Pty) Ltd.
Ground Floor, Block K West, Central Park
400 16th Road, Randjespark,
Midrand,1685
Tel: +27 (0) 11 848 3050

This leaflet was last revised in

28 July 2023

Registration number

Amlodipine 5 Biotech: 37/7.1/0378
Amlodipine 10 Biotech: 37/7.1/0379

AMLODIPINE BIOTECH

PASIËNTINLIGTINGSPAMFLET

SKEDULERINGSTATUS:

53

AMLODIPINE 5 BIOTECH tablette
AMLODIPINE 10 BIOTECH tablette
Amlopidienmaleaat
Suikervry

Lees die hele pamflet versigtig voordat jy begin om AMLODIPINE BIOTECH te neem

- Hou hierdie pamflet. Jy mag dit dalk weer moet lees.
- Indien jy verdere vrae het, vra asseblief jou dokter, apteker, verpleegster of ander gesondheidsorgverskaffer.
- AMLODIPINE BIOTECH is vir jou persoonlik voorgeskryf en jy moet nie jou medisyne met ander mense deel nie. Dit kan hulle skade aandoen, al is hul simptome dieselfde as joune.

Wat is in hierdie pamflet

- Wat AMLODIPINE BIOTECH is em waarvoor dit gebruik word
- Wat jy moet weet voordat jy AMLODIPINE BIOTECH neem
- Hoe om AMLODIPINE BIOTECH te neem
- Moontlike nuwe-effekte
- Hoe om AMLODIPINE BIOTECH te bewaar
- Inhoud van die pak en ander inligting

1. Wat AMLODIPINE BIOTECH is en waarvoor dit gebruik word

Amlopidienmaleaat, die aktiewe bestanddeel van AMLODIPINE 5 BIOTECH en AMLODIPINE 10 BIOTECH, is ’n selektiewe kalsiumkanaal-blokker wat ’n effek het op die bloedvate.

AMLODIPINE BIOTECH word gebruik om angina pectoris en ligte tot matige hipertensie te behandel, alleen of in kombinasie met ander antihipertensiewe medisyne.

2. Wat jy moet weet voordat jy AMLODIPINE BIOTECH neem

Moenie AMLODIPINE BIOTECH neem nie:

- indien jy hipersensitief (allergies) is vir amlopidienmaleaat of vir enige van die ander bestanddele van AMLODIPINE BIOTECH (gelys in afdeling 6).
- indien jy hipersensitief is vir dihidropiridiene.

Waarskuwings en voorsorgmaatreëls

Spesiale sorg moet geneem word met AMLODIPINE BIOTECH:

- indien jy bejaard is en jou dosis verhoog moet word
- indien jy lewersiekte het
- indien jy hartversaking het
- indien jy onlangs ’n hartaanval gehad het

Kinders

Veiligheid en doeltreffendheid is nie vasgestel nie.

Ander medisyne en AMLODIPINE BIOTECH

Vertel altyd jou gesondheidsorgverskaffer indien jy enige ander medisyne neem. (Dit sluit alle komplementêre of tradisionele medisyne in.)

AMLODIPINE BIOTECH kan geaffekteer word deur ander medisyne soos:

- ritonavir, indinavir, nelfinavir (sogenaamde protease-inhibeerders wat gebruik word om MIV te behandel)
- ketokonasool, itrakonasool (teenswammemedisyne)
- rifampisien, eritromisien, klaritromisien (antibiotika)
- verapamil, diltiasem (hartmedisyne)
- hypericum perforatum (St. John Wort)
- dantroleen (infusie vir ernstige liggaamstemperatuurabnormaliteite)

AMLODIPINE BIOTECH mag ander medisyne affekteer, soos:

- takrolimus, sirolimus, temsrolimus, en everolimus (medisyne wat gebruik om die manier waarop jou immuunstelsel werk te verander)
- siklosporien (’n immuunonderdrukker)
- simvastatine (’n cholesterolverlagende medisyne)

AMLODIPINE BIOTECH met kos en drank

Pomelosap en pomelos moet nie ingeneem word deur mense wat AMLODIPINE BIOTECH neem nie. Dit is omdat pomelo en pomelosap kan lei tot ’n toename in die bloedvlakke van die aktiewe bestanddeel amlopidien, wat ’n onvoorspelbare toename in die bloeddrukverlagende effek van amlopidien kan veroorsaak.

Swangerskap, borsvoeding en vrugbaarheid

Indien jy swanger is of borsvoed, dink dat jy dalk mag swanger wees of beplan om ’n baba te hê, raadpleeg asseblief jou dokter, apteker of ander gesondheidsorgverskaffer vir advies voordat jy hierdie medisyne neem.

Swangerskap

Die veiligheid van AMLODIPINE BIOTECH in menslike swangerskap is nog nie vasgestel nie.

Borsvoeding

Amlopidien is bewys dat dit uitgeskei word in klein hoeveelhede in die borsmelk.

Bestuur en die gebruik van masjiene

As jy dalk naar, duiselig, moeg voel of hoofpyn het, moenie bestuur of masjinerie gebruik nie, veral aan die begin van die gebruik van AMLODIPINE BIOTECH.

Dit is nie altyd moontlik om te voorspel tot watter mate AMLODIPINE BIOTECH met jou daaglikse aktiwiteite kan inmeng nie. Jy moet verseker dat jy nie ’n voertuig bestuur of masjiene gebruik totdat jy bewus is van die mate waarin AMLODIPINE BIOTECH jou affekteer nie.

3. Hoe om AMLODIPINE BIOTECH te neem

Moenie medisyne wat vir jou voorgeskryf is met enige ander persoon deel nie.

Neem altyd AMLODIPINE BIOTECH presies soos jou dokter of apteker jou vertel het. Raadpleeg jou dokter of apteker indien jy nie seker is nie.

Jou dokter sal jou vertel hoe lank jou behandeling met AMLODIPINE BIOTECH sal aanhou. Moenie die behandeling vroeg stop nie. Indien jy die indruk het dat die effek van AMLODIPINE BIOTECH te sterk of te swak is, vertel jou dokter of apteker.

Neem die tablette met genoegsame vloeistof (bv. een glas water). Die aanbeveelde dosis van AMLODIPINE BIOTECH is 5 mg per dag. Indien daar geen verbetering is na 10 tot 14 dae van behandeling nie kan AMLODIPINE BIOTECH dosisse verhoog word tot 10 mg een keer per dag.

Indien jy meer AMLODIPINE BIOTECH neem as wat jy moet

Die neem van te veel tablette kan veroorsaak dat jou bloeddruk laag of selfs gevaarlik laag word. Jy kan duiselig, lighoofdig, flou of swak voel. As bloeddrukdaling ernstig is, kan skok voorkom. Jou vel kan koel en klam voel en jy kan jou bewussyn verloor. ’n Oormaat vloeistof kan in jou longe oophou (pulmonêre eeem) wat kortasem veroorsaak wat tot 24-48 uur na inname kan ontwikkel.

In die geval van ’n oordosis, raadpleeg jou dokter of apteker. Indien hulle nie beskikbaar is nie, kontak die naaste hospitaal of gifsentrum.

Indien jy vergeet om AMLODIPINE BIOTECH te neem

Moenie ’n dubbele dosis neem om op te maak vir vergete individuele dosisse nie.

4. Moontlike nuwe-effekte

AMLODIPINE BIOTECH kan nuwe-effekte hê.

Nie alle nuwe-effekte wat vir AMLODIPINE BIOTECH aangemeld is word in hierdie pamflet ingesluit nie. Sou jou algemene gesondheid versleg of indien jy enige ongewensde effekte ervaar terwyl jy AMLODIPINE BIOTECH neem, raadpleeg asseblief jou gesondheidsorgverskaffer vir advies.

Indien enige van die volgende gebeur, hou op om AMLODIPINE BIOTECH te neem/ gebruik en vertel onmiddelik jou dokter of gaan na die ongevalle-afdeling van jou naaste hospitaal:

- swelling van die hande, voete, enkels, lippe en mond of keel wat moeilike asemhaling of sluk veroorsaak,
- veluitslag of jeuk,
- floute.

Hierdie is almal baie ernstige nuwe-effekte. Indien jy hulle het mag jy dalk ’n ernstige reaksie op AMLODIPINE BIOTECH gehad het. Jy mag dalk dringende mediese aandaag of hospitalisasie benodig.

Vertel jou dokter onmiddelik of gaan na die ongevalle-afdeling van jou naaste hospitaal indien jy enige van die volgende opmerk:

- tekort aan bloedplaatjies in die bloed, wat bloeding in die weefsels veroorsaak, kneusing en stadige bloedstolling na besering (trombositopenie)
- vermindering in die aantal wit selle in die bloed (leukositopenie)
- oormaat van glukose in die bloedstroom (hiperglukemie)
- tydelike verlies van bewussyn wat veroorsaak word deur ’n daling in bloeddruk
- vinnig, sterk of onreëlmatige hartklop
- hartaanval
- hartklop is onreëlmatig (stadig, vining) of jy het ’n abnormale ritme (hart se twee boonste kamers (die atria) klop chaoties en onreëlmatig (atriale fibrillasie)
- moeilike asemhaling
- inflammasie van die pankreas
- Inflammasie van die lewer
- Verhoogde lewerensiemer en lewersiekte

Hierdie is almal ernstige nuwe-effekte. Jy mag dalk dringende mediese aandaag benodig.

Vertel jou dokter indien jy enige van die volgende opmerk:

Gereelde nuwe-effekte:

- duiseligheid
- hoofpyn (veral in die begin van behandeling)
- lomerigheid
- visieversteurings en dubbel visie
- blosing, word rooi en warm
- naarheid
- maagpyn
- slegte spysvertering (dispepsie)
- diarree
- hardlywigheid
- swelling van die enkels
- spierkrampe
- uiterse moegheid
- fisiese swakheid of tekort aan energie
- oormatige waterige vloeistof wat in die holtes of weefsel van die liggaam versamel (edeem)

Minder gereelde nuwe-effekte:

- Depressie
- gemoedsveranderinge insluitend gevoel van bekommernis of senuweeaagtigheid
- slaperigheid
- verwarring
- hoë vlakke van spiertonus of spanning (hipertonie)
- verlies van sensasie in die liggaam (hipoestesia)
- naalde en spelde gevoel (parestesia)
- swakheid, gevoelloosheid en pyn, gewoonlik in jou hande en voete (perifere neuropatie)
- onwillekeurige beweging
- disfunksie in die sin van smaak (disgeusia)
- lui of gons in die ore
- lae bloeddruk
- inflammasie van die bloedvate
- hoes
- rinitis
- braking
- droë mond
- oorgroei van tandvleisweefsel rondom die tande
- inflammasie van die voering van die maag
- vergeling van die vel of die wit van die oë
- algehele afwesigheid van hare van areas van die liggaam waar dit normaalweg groei, kaalheid (alopesia)
- ’n veluitslag van pers kolle op die vel wat veroorsaak word deur interne bloeding van klein bloedvate (purpura)
- velverkleuring
- abnormale oormatige sweet wat nie noodwendig verwant is aan hitte of oefening nie (hiperhidrosis)
- jeuk van die vel (pruritus)
- veluitslag
- rooi, verheve, jeukerige veluitslag
- swelling rondom die oë en lippe (angioedeem)
- vel immuunreaksie (erythema multiforme)
- rooiheid en afskliering van die vel (eksfoliatiewe dermatitis)
- is ’n seldsame, ernstige afwyking van die vel en slymvliese. Dit is gewoonlik ’n reaksie op medikasie wat begin met griepagtige simptome, gevolg deur ’n pylnike uitslag wat versprei en blase. Dan sterf die boonste laag van die aangepaste vel, val af en begin na ’n paar dae genees (Steven Johnson se sindroom).
- gewrigspyn
- rugpyn
- spierpyn
- moet meer gereeld as gewoonlik urineer
- abnormaliteite in die proses van urinering, insluitend blaasbeheer, frekwensie van urinering, sowel as die volume en samestelling van urine
- word gedurende die nag wakker om te urineer (nokturie)
- vergroting van ’n man se borste (ginekomasie)
- onvermoë vir ’n man om ’n ereksie of orgasme te verkry (impotensie)
- pyn
- borspyn
- algemene gevoel van ongemak, siekte of ongemak (malaise)
- gewigstoename
- gewigswertes

Nuwe-effekte met onbekende frekwensie:

lewensbedreigende velversteuring gekenmerk deur blasies en afskliering van die vel (toksiese epidermale nekrolise). Indien jy enige nuwe-effekte opmerk wat nie in hierdie pamflet genoem word nie, lig asseblief jou dokter of apteker daarvan in.

Aanmelding van nuwe-effekte

Indien jy nuwe-effekte kry, praat met jou dokter, apteker of verpleegster. Jy kan ook nuwe-effekte aan SAHPRA aanmeld deur die vorm **“6.04 Adverse Drug Reaction Reporting Form”**, wat aanlyn onder SAHPRA se publikasies gevind kan word: https://www.sahpra.org.za/Publications/Index/8. Deur nuwe-effekte aan te meld kan jy help om meer inligting oor die veiligheid van AMLODIPINE BIOTECH te verskaf.

5. Hoe om AMLODIPINE BIOTECH te bewaar

Bewaar alle medisyne buite bereik van kinders. Bewaar teen of benede 25 °C. Beskerm teen vog. Gee alle ongebruikte medisyne terug aan jou apteker. Moenie ongebruikte medisyne in dreine of rioolstelsels (bv. toilette) weggooi nie.

6. Inhoud van die pak en ander inligting

Wat AMLODIPINE BIOTECH bevat

Die aktiewe bestanddeel is amlopidienmaleaat gelykstaande aan amlopidien. AMLODIPINE BIOTECH is beskikbaar in twee sterktes wat of 5 mg of 10 mg amlopidien bevat, onderskeidelik. Die ander bestanddele is mikrokristallyne sellulose (Avicel PH 112), watervrye kalsium waterstofosfaat, magnesiumstereaat en natriumsytselglikolaat (Type A) (natriumkarboksimetielstysel).

Hoe AMLODIPINE BIOTECH lyk en die inhoud van die pak

AMLODIPINE 5 BIOTECH tablette is wit tot naaswit, onbedek, langwerpige tablette. AMLODIPINE 10 BIOTECH tablette is wit tot naaswit, onbedek, langwerpige tablette met ’n breeklyn aan die een kant. AMLODIPINE BIOTECH word verpak in wit, ondeurskynende stulpstroeke en wit, ondeurskynende plastiek sekuriteitshouers van 28, 30 en 100 tablette. PE ritssluiting pasiënt gereed pakkies (PRP’s) is beskikbaar in pakgroottes van 28 tablette. Alle verpakkingsgroottes en sterktes sal dalk nie gelyktydig bemark word nie.

Houer van die sertifikaat van registrasie

Biotech Laboratories (Edms.) Bpk.
Grondvloer, Blok K Wes, Central Park
16^{de} Weg 400, Randjespark
Midrand, 1685
Tel: +27 (0) 11 848 3050

Hierdie pamflet is laas hersien

28 Julie 2023

Registrasienommer

Amlopidine 5 Biotech: 37/7.1/0378

Amlopidine 10 Biotech: 37/7.1/0379