

TORASEMIDE BIOTECH

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS:

S3

TORASEMIDE BIOTECH 5 tablets
TORASEMIDE BIOTECH 10 tablets
TORASEMIDE BIOTECH 20 tablets
Torasemide

Contains sugar (lactose monohydrate):
TORASEMIDE BIOTECH 5 contains 79,0 mg, TORASEMIDE BIOTECH 10 contains 158,0 mg, TORASEMIDE BIOTECH 20 contains 316,0 mg lactose monohydrate per tablet.

Read all of this leaflet carefully before you start taking TORASEMIDE BIOTECH tablets.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, please ask your doctor, pharmacist, nurse or other healthcare provider.
- TORASEMIDE BIOTECH has been prescribed for you personally and you should not share your medicine with other people. It may harm them, even if their symptoms are the same as yours.

What is in this leaflet

1. What TORASEMIDE BIOTECH is and what it is used for
2. What you need to know before you take TORASEMIDE BIOTECH
3. How to take TORASEMIDE BIOTECH
4. Possible side effects
5. How to store TORASEMIDE BIOTECH
6. Contents of the pack and other information

1. What TORASEMIDE BIOTECH is and what it is used for

TORASEMIDE BIOTECH contains a medicine called torasemide. This belongs to a group of medicines called 'diuretics' or 'water tablets'.

TORASEMIDE BIOTECH is used to:

- Treat high blood pressure (hypertension).
- Treat swelling caused by too much water in the body (oedema). The tablets work by making you pass more water (urine).

2. What you need to know before you take TORASEMIDE BIOTECH

Do not take TORASEMIDE BIOTECH:

- If you are hypersensitive (allergic) to torasemide or any of the other ingredients of TORASEMIDE BIOTECH (listed in section 6).
- If you are hypersensitive (allergic) to medicines called 'sulphonylureas'. These are used to treat diabetes (high blood sugar). These medicines include chlorpropamide, glibenclamide, glipizide and tolbutamide.
- If you have kidney problems (renal failure) with absence of urine production (anuria).
- If you have liver problems.
- If you have kidney problems (renal failure) with absence of urine production (anuria).
- If you have liver problems.
- If you are pregnant or breastfeeding your baby. See section 'Pregnancy and breastfeeding'.
- If you have a low volume of blood (hypovolaemia).
- If you have low levels of sodium or potassium in your blood (hyponatraemia or hypokalaemia).
- If you have severe disorders of passing water (urinating).
- If you are aged 12 years or younger.

Warnings and precautions

Take special care with TORASEMIDE BIOTECH:

- If you have low levels of potassium or sodium in your blood (shown in blood tests).
 - If you have an enlarged prostate or have difficulty passing water (urinating).
 - If you have gout or high levels of uric acid in your blood.
 - If you have diabetes (diabetes mellitus).
 - If you have low blood pressure (hypotension).
 - If you have heartbeats problems (cardiac dysrhythmia).
- If any of these apply to you, or if you are not sure, talk to your doctor or pharmacist before you take TORASEMIDE BIOTECH.

Children and adolescents

TORASEMIDE BIOTECH is not for use in children and adolescents below 12 years of age.

Other medicines and TORASEMIDE BIOTECH

Always tell your healthcare provider if you are taking any other medicine. (This includes all complementary or traditional medicines.)

Tell your doctor if you are taking any of the following medicines:

- Medicine for high blood pressure called ACE (angiotensin-converting enzyme) inhibitor inhibitors (for e.g. enalapril).
 - Insulin or other medicine used to treat diabetes.
 - Aminoglycoside or cephalosporin antibiotics such as streptomycin, gentamycin, cephalixin and ceftriaxone.
 - Amphotericin B (antifungal medicine).
 - Anti-inflammatory medicines such as indomethacin or ibuprofen.
 - A medicine used for heart failure or irregular heartbeat such as digoxin.
 - Lithium (medicine to stabilise your mood).
 - Medicines containing salicylates such as aspirin or salazopyrin.
 - Theophylline (a medicine used to treat asthma).
 - Steroid medicines, such as hydrocortisone, prednisolone and dexamethasone.
 - Laxatives.
 - Adrenaline (also called epinephrine) or noradrenaline (also called norepinephrine). These are used to treat low blood pressure.
 - Colestyramine, or other 'ion exchange resins' (used to treat high levels of cholesterol in your blood).
 - Medicines to relax your muscles (muscle relaxants).
 - Cisplatin (used to treat cancer).
 - Probenecid (used to treat gout).
- Dosage adjustment may be required when using blood thinning tablets like warfarin in combination with TORASEMIDE BIOTECH.

TORASEMIDE BIOTECH with food and drink

TORASEMIDE BIOTECH can be taken with or without food.

Pregnancy, breastfeeding and fertility

TORASEMIDE BIOTECH should not be used by pregnant or breastfeeding women.

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other healthcare provider for advice before taking TORASEMIDE BIOTECH.

Driving and using machines

TORASEMIDE BIOTECH may affect your ability to drive a motor vehicle or operate machinery. If you are feeling nauseous, dizzy or weak do not drive a vehicle or operate machinery.

TORASEMIDE BIOTECH contains lactose monohydrate

TORASEMIDE BIOTECH contains lactose monohydrate which is a type of sugar. If you have been told by your doctor that you have an intolerance to some sugars contact your doctor before taking TORASEMIDE BIOTECH.

3. How to take TORASEMIDE BIOTECH

Do not share medicines prescribed for you with any other person. Always take TORASEMIDE BIOTECH exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

High blood pressure

The usual dose for adults and elderly people is between 2,5 mg and 5 mg TORASEMIDE BIOTECH once a day.

Water retention (oedema)

The treatment starts with 5 mg TORASEMIDE BIOTECH once a day. Your doctor or healthcare provider may increase the dosage up to 40 mg TORASEMIDE BIOTECH per day.

If you have the impression that the effect of TORASEMIDE BIOTECH is too strong or too weak, talk to your doctor or pharmacist.

If you take more TORASEMIDE BIOTECH than you should

In the event of an overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison centre as soon as possible. Take this leaflet and any remaining tablets with you, so that the doctor knows what you have taken.

If you forget to take TORASEMIDE BIOTECH

If you forget to take a dose, take it as soon as you remember. Do not take a double or larger dose to make up for the forgotten individual doses. Continue to take the next tablet at the usual time.

4. Possible side effects

TORASEMIDE BIOTECH can have side effects. Not all side effects reported for TORASEMIDE BIOTECH are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking TORASEMIDE BIOTECH, please consult your healthcare provider for advice.

If any of the following happens, stop taking TORASEMIDE BIOTECH and tell your doctor immediately or go to the casualty department at your nearest hospital:

- Swelling of the hands, feet, ankles, face, lips, mouth or throat, which may cause difficulty in swallowing or breathing.
- Rash or itching.
- Fainting.

These are all very serious side effects. If you have them, you may have had a serious reaction to TORASEMIDE BIOTECH. You may need urgent medical attention or hospitalisation.

Tell your doctor immediately or go to the casualty department at your nearest hospital if you notice any of the following:

- Bleeding or bruising more easily than normal.
- Frequent infections such as fever, severe chills, sore throat or mouth ulcers.
- Tiredness, being short of breath and looking pale.
- Blockage of a blood vessel by a blood clot elsewhere in the body.
- Chest pain.
- Increase in heart rate.
- Dizziness and light-headedness.
- Yellowing of the eyes or skin, stomach pain, dark coloured urine.
- Widespread rash with blistering and peeling of large areas of skin, particularly around the mouth, nose, eyes and genitals (Stevens-Johnson syndrome), and a more severe form, causing extensive peeling of the skin (more than 30 % of the body surface – toxic epidermal necrolysis).
- Difficulty in passing urine, increased urge to urinate or more-frequent urination.

Tell your doctor if you notice any of the following:

Frequent side effects:

- Lack of energy, drowsiness, weakness, muscle cramps, restlessness, dehydration, dry mouth and thirst – these may be signs of a disturbance in the balance of electrolytes (salts) in your blood.
 - Headache, dizziness.
 - Loss of appetite, upper stomach pain, nausea (feeling sick), vomiting (being sick), diarrhoea, constipation.
 - Muscle spasms.
 - Fatigue, unusual tiredness or weakness.
- Less frequent side effects:*
- Skin sensitive to the sun.
 - Buzzing or ringing in ears, loss of hearing.
 - Low levels of potassium in your blood. This may cause, feeling weak, feeling sleepy, feeling confused or cramps.
 - High uric acid levels in your blood.
 - Increased blood sugar levels.
 - Increased levels of cholesterol and triglycerides (fats) in your blood.

Frequency unknown side effects

- Blurred or disturbed vision.
 - Confusion.
 - Tingling or numbness of the hands or feet.
 - Spinning sensation, difficulty speaking, slurred speech, loss of co-ordination.
 - Inflammation of the pancreas, with severe upper stomach pain often with nausea (feeling sick) and vomiting (being sick).
- If you notice any other effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects:

If you get side effects, talk to your doctor or pharmacist. You can also report side effects to SAHPRA via the "**6.04 Adverse Drug Reaction Reporting Form**", found online under SAHPRA's publications: <https://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of TORASEMIDE BIOTECH.

5. How to store TORASEMIDE BIOTECH

- Store at or below 25 °C.
- Keep tablets in the outer carton until required for use.
- Keep container well closed.
- STORE ALL MEDICINES OUT OF REACH OF CHILDREN.
- Do not use the tablets after the expiry date printed on the container or blister.
- Return all unused medicine to your pharmacist.
- Do not dispose of unused medicine in drains and sewerage systems (e.g. toilets).

6. Contents of the pack and other information

What TORASEMIDE BIOTECH contains

The active substance is torasemide.
TORASEMIDE BIOTECH 5: Each tablet contains 5 mg torasemide.
TORASEMIDE BIOTECH 10: Each tablet contains 10 mg torasemide.
TORASEMIDE BIOTECH 20: Each tablet contains 20 mg torasemide.
The inactive ingredients are croscopolvidone, lactose monohydrate, magnesium stearate, microcrystalline cellulose and povidone

What TORASEMIDE BIOTECH looks like and contents of the pack

TORASEMIDE BIOTECH 5: White to off white, oval shaped, scored tablets, debossed with '5' on plain side and scored on the other side.
TORASEMIDE BIOTECH 10: White to off white, oval shaped, scored tablets, debossed with '10' on plain side and scored on the other side.
TORASEMIDE BIOTECH 20: White to off white, oval shaped, scored tablets, debossed with '20' on plain side and scored on the other side.

TORASEMIDE BIOTECH 5 and 10:

30 and 100 tablets are packed in white HDPE container with white ribbed child resistant HDPE cap with a pulp liner in between the heat seal liner and inner cap, placed in outer cardboard carton.

TORASEMIDE BIOTECH 20:

100 tablets are packed in white HDPE container with white ribbed child resistant HDPE cap with a pulp liner in between the heat seal liner and inner cap, placed in outer cardboard carton.
All pack size and strengths may not necessarily be marketed at one time.

Holder of certificate of registration

Biotech Laboratories (Pty) Ltd
Ground Floor, Block K West, Central Park
400 16th Road, Randjespark
Halfway House
Midrand 1685
011 848 3050

This leaflet was last revised

Date of registration: 26 November 2015
Date of revision of the text: 05 February 2024

Registration numbers

TORASEMIDE BIOTECH 5: 45/18.1/1185
TORASEMIDE BIOTECH 10: 45/18.1/1186
TORASEMIDE BIOTECH 20: 45/18.1/1187

TORASEMIDE BIOTECH

PASIËNTINLIGTINGSPAMFLET

SKEDULERINGSTATUS:

[53]

TORASEMIDE BIOTECH 5 tablette
TORASEMIDE BIOTECH 10 tablette
TORASEMIDE BIOTECH 20 tablette
Torasemied

Bevat suiker (laktosemonohidraat):
TORASEMIDE BIOTECH 5 bevat 79,0 mg, TORASEMIDE BIOTECH 10 bevat 158,0 mg, TORASEMIDE BIOTECH 20 bevat 316,0 mg laktosemonohidraat per tablet.

Lees die hele pamflet noukeurig deur voordat jy begin om TORASEMIDE BIOTECH tablette te neem.

- Hou hierdie pamflet. Jy mag dit dalk weer moet deurlees.
- Indien jy enige verdere vrae het, vra asseblief jou dokter, apteker, verpleegster of ander gesondheidsorgverskaffer.
- TORASEMIDE BIOTECH is vir jou persoonlik voorgeskryf en jy moet nie jou medisyne met ander mense deel nie. Dit kan hulle benadeel, selfs al is hul simptome dieselfde as joune.

Wat is in hierdie pamflet

- Wat TORASEMIDE BIOTECH is en waarvoor dit gebruik word
- Wat jy moet weet voordat jy TORASEMIDE BIOTECH neem
- Hoe om TORASEMIDE BIOTECH te neem
- Moontlike newe-effekte
- Hoe om TORASEMIDE BIOTECH te bewaar
- Inhoud van die pak en ander inligting

1. Wat TORASEMIDE BIOTECH is en waarvoor dit gebruik word
TORASEMIDE BIOTECH bevat ’n medisyne genaamd torasemied. Dit behoort aan ’n groep medisyne wat ‘diuretika’ of ‘watertablette’ genoem word.

TORASEMIDE BIOTECH word gebruik om:

- Behandel hoë bloeddruk (hipertensie).
- Behandel swelling wat veroorsaak word deur te veel water in die liggaam (edeem).

Die tablette werk deur jou meer water (uriene) te laat uitlaat.

2. Wat jy moet weet voordat jy TORASEMIDE BIOTECH neem
Moenie TORASEMIDE BIOTECH neem nie:

- Indien jy hipersensitief (allergies) is vir torasemied of enige van die ander bestanddele van TORASEMIDE BIOTECH (gelys in afdeling 6).
- Indien jy hipersensitief (allergies) is vir medisyne genaamd ‘sulfonylureums’. Dit word gebruik om diabetes (hoë bloedsuiker) te behandel. Hierdie medisyne sluit in chloorpropamied, glibenklamied, glipisied en tolbutamied.
- Indien jy nierprobleme (nierversaking) het met afwesigheid van urieneproduksie (anurie).
- Indien jy lewerprobleme het.
- Indien jy swanger is of jou baba borsvoed. Sien afdeling ‘Swangerskap en borsvoeding’.
- Indien jy ’n lae volume bloed het (hipovolemie).
- Indien jy lae vlakke van natrium of kalium in jou bloed het (hiponatremie of hipokalemie).
- Indien jy ernstige afwykings van deurvloeiende water het (urineer).
- Indien jy 12 jaar of jonger is.

Waarswakings en voorsorgmaatreëls

Wees vaal versigtig met TORASEMIDE BIOTECH:

- Indien jy lae vlakke van kalium of natrium in jou bloed het (getoet in bloedtoetse).
- Indien jy ’n vergrote prostaat het of sukkel om water deur te gee (urineer).
- Indien jy jig of hoë vlakke van uriensuur in jou bloed het.
- Indien jy diabetes/ suikersiekte (diabetes mellitus) het.
- Indien jy lae bloeddruk (hipotensie) het.
- Indien jy hartklopprobleme het (kardiale disritmie).

Indien enige van hierdie op jou van toepassing is, of as jy nie seker is nie, praat met jou dokter of apteker voordat jy TORASEMIDE BIOTECH neem.

Kinders en adolessente

TORASEMIDE BIOTECH is nie vir gebruik by kinders en adolessente jonger as 12 jaar nie.

Ander medisyne en TORASEMIDE BIOTECH

Vertel altyd jou gesondheidsorgverskaffer indien jy enige ander medisyne gebruik. (Dit sluit alle komplementêre of tradisionele medisyne in.)

Vertel jou dokter indien jy enige van die volgende medisyne gebruik:

- Medisyne vir hoë bloeddruk genoem ACE (angiotensien-omskakelende ensiem) inhibeerders (bv. enalapril).
- Insulien of ander medisyne wat gebruik word om diabetes te behandel.
- Aminoglikosied- of kefalosporien-antibiotika soos streptomisien, gentamisien, kefaleksien en seftriaksoon.
- Amfoterisien B (teenswam medisyne).
- Anti-inflammatoriese medisyne soos indometasien of ibuprofen.
- ’n Medisyne wat gebruik word vir hartversaking of onreëlmatige hartklop soos digoksiein.
- Litium (medisyne om jou bui te stabiliseer).
- Medisyne wat salisilate bevat soos aspirien of salasopirien.
- Teofillien (’n medisyne wat gebruik word om asma te behandel).
- Steroïedmedisyne, soos hidrokortisoon, prednisoloon en deksametasoon.
- Lakseermiddels.
- Adrenaliën (ook genoem epinefrien) of noradrenalin (ook genoem norepinefrien). Dit word gebruik om lae bloeddruk te behandel.
- Kolestriamien, of ander ‘ioonuitruilharse’ (word gebruik om hoë vlakke van cholesterol in jou bloed te behandel).
- Medisyne om jou spiere te ontspan (spierverslappers).
- Cisplatin (gebruik om kanker te behandel).
- Probenesied (gebruik om jig te behandel).

Dosisaanpassing mag nodig wees wanneer bloedverdundingstablette soos warfarin in kombinasie met TORASEMIDE BIOTECH gebruik word.

TORASEMIDE BIOTECH met kos en drank

TORASEMIDE BIOTECH kan met of sonder kos geneem word.

Swangerskap, borsvoeding en vrugbaarheid

TORASEMIDE BIOTECH moet nie deur swanger of borsvoedende vroue gebruik word nie.

Indien jy swanger is of borsvoed, dink jy is dalk swanger of beplan om ’n baba te hê, raadpleeg asseblief jou dokter, apteker of ander gesondheidsorgverskaffer vir advies voordat TORASEMIDE BIOTECH geneem word.

Bestuur en gebruik van masjiene

TORASEMIDE BIOTECH kan jou vermoë om ’n motorvoertuig te bestuur of masjinerie te bestuur, beïnvloed. As jy naar, duiselig of swak voel, moenie ’n voertuig bestuur of masjinerie gebruik nie.

TORASEMIDE BIOTECH bevat laktosemonohidraat

TORASEMIDE BIOTECH bevat laktosemonohidraat wat ’n soort suiker is. Indien jy deur jou dokter gesê het dat jy ’n onverdraagsaamheid teenoor sommige suikers het, kontak jou dokter voordat jy TORASEMIDE BIOTECH neem.

3. Hoe om TORASEMIDE BIOTECH te neem

Moenie medisyne wat vir jou voorgeskryf is met enige ander persoon deel nie.

Neem altyd TORASEMIDE BIOTECH presies soos jou dokter vir jou gesê het. Gaan met jou dokter of apteker as jy nie seker is nie.

Hoë bloeddruk

Die gewone dosis vir volwassenes en bejaardes is tussen 2,5 mg en 5 mg TORASEMIDE BIOTECH een keer per dag.

Waterretensie (edeem)

Die behandeling begin met 5 mg TORASEMIDE BIOTECH een keer per dag. Jou dokter of gesondheidsorgverskaffer kan die dosis verhoog tot 40 mg TORASEMIDE BIOTECH per dag.

Indien jy die indruk het dat die effek van TORASEMIDE BIOTECH te sterk of te swak is, praat met jou dokter of apteker.

Indien jy meer TORASEMIDE BIOTECH neem as wat jy moet

In die geval van ’n oordosis, raadpleeg jou dokter of apteker. Indien nie een beskikbaar is nie, kontak die naaste hospitaal of gifsentrum so gou moontlik. Neem hierdie pamflet en enige orbylvende tablette saam, sodat die dokter weet wat jy geneem het.

Indien jy vergeet om TORASEMIDE BIOTECH te neem

As jy vergeet om ’n dosis te neem, neem dit so gou as wat jy onthou. Moenie ’n dubbele of groter dosis neem om op te maak vir die vergeete individuele dosisse nie. Gaan voort om die volgende tablet op die gewone tyd te neem.

4. Moontlike newe-effekte

TORASEMIDE BIOTECH kan newe-effekte hê. Nie alle newe-effekte wat vir TORASEMIDE BIOTECH aangemeld is, is in hierdie pamflet ingesluit nie. Indien jou algemene gesondheid versleg of as jy enige nadelige effekte ervaar terwyl jy TORASEMIDE BIOTECH neem, raadpleeg asseblief jou gesondheidsorgverskaffer vir advies.

Indien enige van die volgende gebeur, hou op om TORASEMIDE BIOTECH te neem en vertel jou dokter dadelik of gaan na die ongevalle-afdeling by jou naaste hospitaal:

- Swelling van die hande, voete, enkels, gesig, lippe, mond of keel, wat probleme mkan veroorsaak om te sluk of asem te haal.
- Uitslag of jeuk.
- Floute.

Hierdie is alles baie ernstige newe-effekte. As jy dit het, het jy dalk ’n ernstige reaksie op TORASEMIDE BIOTECH gehad. Jy benodig dalk dringende mediese aandag of hospitalisasie.

Vertel jou dokter dadelik of gaan na die ongevalle-afdeling by jou naaste hospitaal indien jy enige van die volgende opmerk:

- Bloei of kneusing makliker as normaal.
- Gereelde infeksies soos koors, erg kouekoors, seer keel of mondseer.
- Moegheid, kortasem en bleek lyk.
- Blokkering van ’n bloedvat deur ’n bloedklont elders in die liggaam.
- Borspyn.
- Verhoging in hartklop.
- Duiseligheid en lighoofdigheid.
- Vergeling van die oë of vel, maagpyn, donkerkleurige uriene.
- Wydverspreide uitslag met blase en afskifering van groot dele van die vel, veral rondom die mond, neus, oë en geslagsdele (Stevens-Johnson-sindroom), en ’n meer ernstige vorm, wat uitgebreide afskifering van die vel veroorsaak (meer as 30 % van die liggaam) oppervlak – toksiese epidermale nekrolise).
- Moelijk urineer, verhoogde drang om te urineer of meer gereelde urineering.

Vertel jou dokter indien jy enige van die volgende opmerk:

Gereelde newe-effekte:

- Gebreke aan energie, lomerigheid, swakheid, spierkrampe, rusteloosheid, dehidrasie, droë mond en dors – dit kan tekens wees van ’n versteuring in die balans van elektroliete (soute) in jou bloed.
- Hoofpyn, duiseligheid.
- Verlies aan eetlust, boonste maagpyn, naarheid (siek voel), braking (siek wees), diarree, hardlywigheid.
- Spierspasmas.
- Moegheid, ongewone moegheid of swakheid.

Minder gereelde newe-effekte:

- Vel sensitief vir die son.
- Gezoem of gelui in ore, gehoorverlies.
- Lae vlakke van kalium in jou bloed. Dit kan veroorsaak, swak voel, slaperig voel, verward of krampe voel.
- Hoë uriensuurvlakke in jou bloed.
- Verhoogde bloedsuikervlakke.
- Verhoogde vlakke van cholesterol en trigliseriede (vette) in jou bloed.

Newe-effekte met onbekende frekwensie:

- Dowwe of versteurde visie.
- Verwarring.
- Tinteling of gevoelloosheid van die hande of voete.
- Spin-senssie, probleme met praat, onduidelike spraak, verlies aan koördinasie.
- Inflammasie van die pankreas, met erge boonste maagpyn dikwels met naarheid (siek voel) en braking (siek wees).

Indien jy enige ander effekte opmerk wat nie in hierdie pamflet genoem word nie, stel asseblief jou dokter of apteker in kennis.

Aanmelding van newe-effekte:

Indien jy newe-effekte kry, praat met jou dokter of apteker. Jy kan ook newe-effekte aan SAHPRA aanmeld deur die vorm **“6.04 Adverse Drug Reaction Reporting Form”**, wat aanlyn onder SAHPRA se publiskasies gevind kan word: https://www.sahpra.org.za/Publications/Index/8. Deur newe-effekte aan te meld kan jy help om meer inligting oor die veiligheid van TORASEMIDE BIOTECH te verskaf.

5. Hoe om TORASEMIDE BIOTECH te bewaar

- Bewaar teen of benede 25 °C.
- Hou tablette in die buitekarton totdat dit benodig word vir gebruik.
- Hou houer goed toe.
- BEWAAR ALLE MEDISYNE BUITE BEREIK VAN KINDERS.**
- Moenie die tablette na die vervaldatum wat op die houer of stulpstrok gedruk is, gebruik nie.
- Geel alle ongebruikte medisyne terug aan jou apteker.
- Moenie ongebruikte medisyne in dreine en rioolstelsels (bv. toilette) weggooi nie.

6. Inhoud van die pak en ander inligting
Wat TORASEMIDE BIOTECH bevat

Die aktiewe stof is torasemied.

TORASEMIDE BIOTECH 5: Elke tablet bevat 5 mg torasemied.
TORASEMIDE BIOTECH 10: Elke tablet bevat 10 mg torasemied.
TORASEMIDE BIOTECH 20: Elke tablet bevat 20 mg torasemied.
Die onaktiewe bestanddele is krosprovidoon, laktosemonohidraat, magnesiumstearaat, mikrokristallyne selulose en povidoon

Hoe TORASEMIDE BIOTECH lyk en die inhoud van die pak
TORASEMIDE BIOTECH 5: Wit tot naaswit, ovaalvormige, tablette met ’n kerfstreep, ingedruk met ‘5’ op die gewone kant en met ’n kerf aan die ander kant.

TORASEMIDE BIOTECH 10: Wit tot naaswit, ovaalvormige tablette met kerfgate, gegraveer met ‘10’ op die gewone kant en gekerf aan die ander kant.

TORASEMIDE BIOTECH 20: Wit tot naaswit, ovaalvormige tablette met ’n kerfstreep, gegraveer met ‘20’ op die gewone kant en gekerf aan die ander kant.

TORASEMIDE BIOTECH 5 en 10:

30 en 100 tablette is verpak in ’n wit HDPE-houer met ’n wit geribbelde kinderbestande HDPE-dop met ’n pulpvoering tussen die hitteseëlvoering en binnedop, geplaas in buitenste kartonhouer.
TORASEMIDE BIOTECH 20: 100 tablette is verpak in ’n wit HDPE-houer met ’n wit geribbelde kinderbestande HDPE-dop met ’n pulpvoering tussen die hitteseëlvoering en binnedop, geplaas in buitenste kartonhouer.
Alle verpakkingsgroottes en sterktes mag nie noodwendig op een slag bemark word nie.

Houer van die sertifikaat van registrasie

Biotech Laboratories (Edms) Bpk
Grondvloer, Blok K Wes, Central Park
16^{de} Weg, 400, Randjespark
Halfweghuis
Midrand 1685
011 848 3050

Hierdie pamflet is laas hersien

Datum van registrasie: 26 November 2015

Datum van hersiening van die teks: 05 Februarie 2024

Registrasienuommers

TORASEMIDE BIOTECH 5: 45/18.1/1185

TORASEMIDE BIOTECH 10: 45/18.1/1186

TORASEMIDE BIOTECH 20: 45/18.1/1187